

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed	for						Go	Clear
Limits		Preview/Index		History		Clipboard		Details
Display	Abstract	✓	Show: 20	✓	Sort	✓	Send to	Text

☐ 1: J Gen Microbiol. 1991 Aug;137 (Pt 8):1911-21.

[Related Articles, Links](#)

Entrez
PubMed

Molecular and evolutionary relationships among enteric bacteria.

Lawrence JG, Ochman H, Hartl DL.

Department of Genetics, Washington University School of Medicine, St. Louis, MO 63110.

PubMed
Services

Classification of bacterial species into genera has traditionally relied upon variation in phenotypic characteristics. However, these phenotypes often have a multifactorial genetic basis, making unambiguous taxonomic placement of new species difficult. By designing evolutionarily conserved oligonucleotide primers, it is possible to amplify homologous regions of genes in diverse taxa using the polymerase chain reaction and determine their nucleotide sequences. We have constructed a phylogeny of some enteric bacteria, including five species classified as members of the genus *Escherichia*, based on nucleotide sequence variation at the loci encoding glyceraldehyde-3-phosphate dehydrogenase and outer membrane protein 3A, and compared this genealogy with the relationships inferred by biotyping. The DNA sequences of these genes defined congruent and robust phylogenetic trees indicating that they are an accurate reflection of the evolutionary history of the bacterial species. The five species of *Escherichia* were found to be distantly related and, contrary to their placement in the same genus, do not form a monophyletic group. These data provide a framework which allows the relationships of additional species of enteric bacteria to be inferred. These procedures have general applicability for analysis of the classification, evolution, and epidemiology of bacterial taxa.

Related
Resources

PMID: 1955870 [PubMed - indexed for MEDLINE]

Display	Abstract	✓	Show: 20	✓	Sort	✓	Send to	Text
---------	----------	---	----------	---	------	---	---------	------

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
[Freedom of Information Act](#) | [Disclaimer](#)

Jun 5 2003 10:08:34



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books
Search PubMed for [] Go Clear
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Infect Immun 1994 Oct;62(10):4632-6

[Related Articles, Links](#)

Entrez PubMed

[Overview](#)

[Help | FAQ](#)

[Tutorial](#)

[New/Noteworthy](#)

[E-Utilities](#)

PubMed Services

[Journals Database](#)

[MeSH Browser](#)

[Single Citation Matcher](#)

[Batch Citation Matcher](#)

[Clinical Queries](#)

[LinkOut](#)

[Cubby](#)

Related Resources

[Order Documents](#)

[NLM Gateway](#)

[TOXNET](#)

[Consumer Health](#)

[Clinical Alerts](#)

[ClinicalTrials.gov](#)

[PubMed Central](#)

[Privacy Policy](#)

Biological activities of native and recombinant *Borrelia burgdorferi* outer surface protein A: dependence on lipid modification.

Weis JJ, Ma Y, Erdile LF.

Department of Pathology, University of Utah School of Medicine, Salt Lake City 84132.

Borrelia burgdorferi lipoproteins are 50- to 500-fold more active as cytokine inducers and B-cell mitogens than *Escherichia coli* lipoproteins and synthetic peptides containing the tripalmitoyl-S-glycerol-cysteine moiety. To investigate the source of this unique potency, we compared native OspA from *B. burgdorferi* with recombinant lipidated OspA produced in *E. coli*. As little as 10 ng of either protein per ml stimulated B-cell proliferation and production of cytokines and nitric oxide by macrophages. The two proteins induced comparable antibody responses in mice. Nonlipidated OspA made in *E. coli* had no stimulatory activity. Thus, lipid modification is essential both in vivo and in vitro for the immunological properties of OspA. The lipid moiety appears equally active whether produced in *B. burgdorferi* or in *E. coli*.

PMID: 7927731 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)

[NCBI | NLM | NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act | Disclaimer](#)

i686-pc-linux-gnu Jan 21 2003 17:57:06



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books
Search PubMed for Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Infect Immun 1993 Jan;61(1):81-90

[Related Articles, Links](#)

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

Role of attached lipid in immunogenicity of *Borrelia burgdorferi* OspA.

Erdile LF, Brandt MA, Warakomski DJ, Westrack GJ, Sadziene A, Barbour AG, Mays JP.

Connaught Laboratories, Inc., Swiftwater, Pennsylvania 18370.

OspA is a protective antigen of the Lyme disease spirochete *Borrelia burgdorferi*. Expression of the full-length *B. burgdorferi* B31 OspA gene in *Escherichia coli* produces a protein that is processed posttranslationally by signal peptidase II and contains an attached lipid moiety. The recombinant OspA lipoprotein has been purified by detergent extraction and ion-exchange chromatography. Priming and boosting with OspA lipoprotein, either with no adjuvant or adsorbed to alum, elicited a strong, dose-dependent immunoglobulin G response. Serum from vaccinated mice inhibited spirochetal growth in vitro. Mice immunized twice with as little as 0.4 micrograms of OspA lipoprotein were protected against an intradermal challenge with 10(4) infectious spirochetes. The ability of the purified recombinant lipoprotein to induce a strong protective response in the absence of toxic adjuvants makes it an excellent candidate antigen for a human vaccine against Lyme disease. By contrast, no serum immunoglobulin G or growth inhibitory response to OspA nonlipoprotein was seen at any dose. The difference in immunogenicities of the lipoprotein and nonlipoprotein forms of OspA is not due to any difference in the antigenicities of the two proteins. These results suggest that posttranslational lipid attachment is a critical determinant of the immunogenicity of the OspA protein.

PMID: 8418068 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

i686-pc-linux-gnu Jan 21 2003 17:57:06



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books
Search PubMed for [] Go Clear
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Eur J Biochem 1998 Jul 15;255(2):446-54

[Related Articles, Links](#)

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

[Privacy Policy](#)

Online

The recombinant *Klebsiella pneumoniae* outer membrane protein OmpA has carrier properties for conjugated antigenic peptides.

Haeuw JF, Raully I, Zanna L, Libon C, Andreoni C, Nguyen TN, Baussant T, Bonnefoy JY, Beck A.

Centre d'Immunologie Pierre Fabre, Saint Julien en Genevois, France.
jean.francois.haeuw@pierre-fabre.com

Klebsiella pneumoniae OmpA, the 40-kDa major protein of the outer membrane, was cloned and expressed in *Escherichia coli*. The recombinant protein was produced intracellularly in *E. coli* as inclusion bodies. Fusion of a short peptide to the N-terminus of native P40 facilitated high-level expression of the recombinant protein. Purified recombinant P40 was analyzed to verify purity and structural integrity. The molecular mass of purified recombinant P40 determined by electrospray mass spectrometry was 37,061 Da, in agreement with the theoretical mass deduced from the DNA sequence. Specific proliferation of recombinant-P40-primed murine lymph node cells in response to recombinant P40 stimulation in vitro indicated the presence of a T-cell epitope on recombinant P40. The induction of high serum antibody titers to a synthetic peptide derived from the attachment protein G of the respiratory syncytial virus when chemically coupled to recombinant P40 indicated that the protein had potent carrier properties.

PMID: 9716387 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

i686-pc-linux-gnu Jan 21 2003 17:57:06



Infect. Immun., 10 1994, 4632-4636, Vol 62, No. 10
Copyright © 1994, American Society for Microbiology

Biological activities of native and recombinant *Borrelia burgdorferi* outer surface protein A: dependence on lipid modification

JJ Weis, Y Ma and LF Erdile

Department of Pathology, University of Utah School of Medicine, Salt Lake City 84132.

- ▶ Similar articles found in:
[IAI Online](#)
[PubMed](#)
 - ▶ [PubMed Citation](#)
 - ▶ This Article has been cited by:
[other online articles](#)
 - ▶ Search Medline for articles by:
[Weis, J. J.](#) || [Erdile, L. F.](#)
 - ▶ Alert me when:
[new articles cite this article](#)
 - ▶ [Download to Citation Manager](#)
-
- ▶ [Books from ASM Press](#)

Borrelia burgdorferi lipoproteins are 50- to 500-fold more active as cytokine inducers and B-cell mitogens than *Escherichia coli* lipoproteins and synthetic peptides containing the tripalmitoyl-S- glyceryl-cysteine moiety. To investigate the source of this unique potency, we compared native OspA from *B. burgdorferi* with recombinant lipidated OspA produced in *E. coli*. As little as 10 ng of either protein per ml stimulated B-cell proliferation and production of cytokines and nitric oxide by macrophages. The two proteins induced comparable antibody responses in mice. Nonlipidated OspA made in *E. coli* had no stimulatory activity. Thus, lipid modification is essential both in vivo and in vitro for the immunological properties of OspA. The lipid moiety appears equally active whether produced in *B. burgdorferi* or in *E. coli*.

This article has been cited by other articles:

- Cote-Sierra, J., Bredan, A., Toldos, C. M., Stijlemans, B., Brys, L., Cornelis, P., Segovia, M., de Baetselier, P., Revets, H. (2002). Bacterial Lipoprotein-Based Vaccines Induce Tumor Necrosis Factor-Dependent Type 1 Protective Immunity against *Leishmania major*. *Infect. Immun.* 70: 240-248 [[Abstract](#)] [[Full Text](#)]
- Wooten, R. M., Ma, Y., Yoder, R. A., Brown, J. P., Weis, J. H., Zachary, J. F., Kirschning, C. J., Weis, J. J. (2002). Toll-Like Receptor 2 Is Required for Innate, But Not Acquired, Host Defense to *Borrelia burgdorferi*. *The JI* 168: 348-355 [[Abstract](#)] [[Full Text](#)]
- Dorward, D. W., Larson, R. S. (2001). Murine Model for Lymphocytic Tropism by *Borrelia burgdorferi*. *Infect. Immun.* 69: 1428-1432 [[Abstract](#)] [[Full Text](#)]
- Talkington, J., Nickell, S. P. (2001). Role of Fc Gamma Receptors in Triggering Host Cell Activation and Cytokine Release by *Borrelia burgdorferi*. *Infect. Immun.* 69: 413-419 [[Abstract](#)] [[Full Text](#)]
- Sellati, T. J., Bouis, D. A., Kitchens, R. L., Darveau, R. P., Pugin, J., Ulevitch, R. J., Gangloff, S. C., Goyert, S. M., Norgard, M. V., Radolf, J. D. (1998). *Treponema pallidum* and *Borrelia burgdorferi* Lipoproteins and Synthetic Lipopeptides Activate Monocytic Cells via a CD14-Dependent Pathway Distinct from That Used by Lipopolysaccharide. *The JI* 160: 5455-5464 [[Abstract](#)] [[Full Text](#)]
- Wooten, R. M., Morrison, T. B., Weis, J. H., Wright, S. D., Thieringer, R., Weis, J. J. (1998). The Role of CD14 in Signaling Mediated by Outer Membrane Lipoproteins of *Borrelia burgdorferi*. *The JI* 160: 5485-5492 [[Abstract](#)] [[Full Text](#)]
- Hirschfeld, M., Kirschning, C. J., Schwandner, R., Wesche, H., Weis, J. H., Wooten, R. M., Weis, J. J. (1999). Cutting Edge: Inflammatory Signaling by *Borrelia burgdorferi* Lipoproteins Is Mediated by

Toll-Like Receptor 2. *The JI* 163: 2382-2386 [[Abstract](#)] [[Full Text](#)]

- Soler-Rodriguez, A. M., Zhang, H., Lichenstein, H. S., Qureshi, N., Niesel, D. W., Crowe, S. E., Peterson, J. W., Klimpel, G. R. (2000). Neutrophil Activation by Bacterial Lipoprotein Versus Lipopolysaccharide: Differential Requirements for Serum and CD14. *The JI* 164: 2674-2683 [[Abstract](#)] [[Full Text](#)]
- Udalova, I. A., Vidal, V., Scragg, I. G., Kwiatkowski, D. (2000). Direct Evidence for Involvement of NF-kappa B in Transcriptional Activation of Tumor Necrosis Factor by a Spirochetal Lipoprotein. *Infect. Immun.* 68: 5447-5449 [[Abstract](#)] [[Full Text](#)]
- Scragg, I. G., Kwiatkowski, D., Vidal, V., Reason, A., Paxton, T., Panico, M., Dell, A., Morris, H. (2000). Structural Characterization of the Inflammatory Moiety of a Variable Major Lipoprotein of *Borrelia recurrentis*. *J. Biol. Chem.* 275: 937-941 [[Abstract](#)] [[Full Text](#)]
- Brightbill, H. D., Libraty, D. H., Krutzik, S. R., Yang, R., Belisle, J. T., Bleharski, J. R., Maitland, M., Norgard, M. V., Plevy, S. E., Smale, S. T., Brennan, P. J., Bloom, B. R., Godowski, P. J., Modlin, R. L. (1999). Host Defense Mechanisms Triggered by Microbial Lipoproteins Through Toll-Like Receptors. *Science* 285: 732-736 [[Abstract](#)] [[Full Text](#)]
- Lien, E., Sellati, T. J., Yoshimura, A., Flo, T. H., Rawadi, G., Finberg, R. W., Carroll, J. D., Espevik, T., Ingalls, R. R., Radolf, J. D., Golenbock, D. T. (1999). Toll-like Receptor 2 Functions as a Pattern Recognition Receptor for Diverse Bacterial Products. *J. Biol. Chem.* 274: 33419-33425 [[Abstract](#)] [[Full Text](#)]
- Brown, J. P., Zachary, J. F., Teuscher, C., Weis, J. J., Wooten, R. M. (1999). Dual Role of Interleukin-10 in Murine Lyme Disease: Regulation of Arthritis Severity and Host Defense. *Infect. Immun.* 67: 5142-5150 [[Abstract](#)] [[Full Text](#)]
- Sigal, L. H. (1997). LYME DISEASE: A Review of Aspects of Its Immunology and Immunopathogenesis. *Annu. Rev. Immunol.* 15: 63-92 [[Abstract](#)] [[Full Text](#)]
- Talkington, J., Nickell, S. P. (1999). *Borrelia burgdorferi* Spirochetes Induce Mast Cell Activation and Cytokine Release. *Infect. Immun.* 67: 1107-1115 [[Abstract](#)] [[Full Text](#)]
- Zhang, H., Niesel, D. W., Peterson, J. W., Klimpel, G. R. (1998). Lipoprotein Release by Bacteria: Potential Factor in Bacterial Pathogenesis. *Infect. Immun.* 66: 5196-5201 [[Abstract](#)] [[Full Text](#)]
- Giambartolomei, G. H., Dennis, V. A., Philipp, M. T. (1998). *Borrelia burgdorferi* Stimulates the Production of Interleukin-10 in Peripheral Blood Mononuclear Cells from Uninfected Humans and Rhesus Monkeys. *Infect. Immun.* 66: 2691-2697 [[Abstract](#)] [[Full Text](#)]

[HOME](#) [HELP](#) [FEEDBACK](#) [SUBSCRIPTIONS](#) [ARCHIVE](#) [SEARCH](#)

[J. Bacteriol.](#)

[J. Virol.](#)

[Eukaryot. Cell](#)

[Microbiol. Mol. Bio. Review](#)

[All ASM Journals](#)

Copyright © 1994 by the American Society for Microbiology. All rights reserved.